

Deep-Ocean Crusts as Telescopes: Using Live Radioisotopes to Probe Supernova Nucleosynthesis

Brian D. Fields¹

Center for Theoretical Astrophysics, Department of Astronomy, University of Illinois, Urbana, IL 61801, USA

Kathrin A. Hochmuth

Department of Physics, University of Illinois, Urbana, IL 61801, USA

and

John Ellis

Theory Division, Physics Department, CERN, 1211 Geneva 23, Switzerland

ABSTRACT

Live ^{60}Fe has recently been detected in a deep-ocean ferromanganese crust, isolated in layers dating from about 3 Myr ago. Since ^{60}Fe has a mean life of 2.2 Myr, a near-Earth supernova is the only likely source for such a signal, and we explore here the consequences of a supernova origin. We combine the ^{60}Fe data with several supernova nucleosynthesis models to calculate the supernova distance as a function of progenitor mass, finding an allowed range of 15 – 120 pc. We also predict the signals expected for several other radioisotopes, which are independent of the supernova distance. Species likely to be present near or above background levels are ^{10}Be , ^{26}Al , ^{53}Mn , ^{182}Hf and ^{244}Pu . Of these, ^{182}Hf and ^{244}Pu are nearly background-free, presenting the best opportunities to provide strong confirmation of the supernova origin of the ^{60}Fe signal, and to demonstrate that at least some supernovae are the source for the r -process. The accuracies of our predictions are hampered by large uncertainties in the predicted ^{60}Fe yields for supernovae of different masses, so the new crust data motivate a redoubled theoretical attack on this problem.

Subject headings: supernovae: general — nuclear reactions, nucleosynthesis, abundances — solar neighborhood

¹also Department of Physics, University of Illinois

1. Introduction

The possibility of finding terrestrial signatures of near-earth supernova explosions has been intensively discussed over the last decade (see, e.g., Ellis, Fields, & Schramm 1996; Fields & Ellis 1999). Recent evidence for a supernova remnant 12° in diameter, that may be associated with ^{26}Al γ -rays, points to an explosion within the past few Myr and within 100–200 pc (McCullough, Fields, & Pavlidou 2002). Moreover, the existence of the Local Bubble – containing hot, rarefied gas in which the Sun resides (Frisch 1995) – suggests that we live in the *interior* of one or more supernova remnants (e.g., Smith & Cox 2001). This enhances the prospects of identifying terrestrial evidence of supernova remnant engulfing the Solar System and the Earth.

Even if it were too far away to be hazardous for life on Earth (see, e.g., Ruderman 1975; Ellis & Schramm 1995), a sufficiently close supernova explosion would deposit on the Earth supernova ejecta and related material produced by cosmic rays (Ellis, Fields, & Schramm 1996). This could have resulted in the possible uptake of supernova-produced elements in deep-ocean sediments or manganese-iron crusts. Following Ellis, Fields, & Schramm (1996), Knie et al. (1999) discovered the first evidence for live ^{60}Fe in a deep-ocean ferromanganese crust, at a level ~ 100 times the expected background and consistent with an event in the last ~ 5 Myr at a distance ~ 30 pc (Fields & Ellis 1999). However, the structure and geometry of the crust made it difficult to reconstruct a reliable time history, and thus the nature of the ^{60}Fe signal remained uncertain.

Very recently, Knie et al (2004) have studied a different ferromanganese crust, which is better suited for a high-precision study with good time resolution. They find a ^{60}Fe spike far above background in a crust layer which formed between 2.4 and 3.2 Myr ago. This detection confirms the earlier result, but now goes beyond detecting the presence of ^{60}Fe to revealing the details of its deposition history. The presence of a live radioisotopic signal in a single, isolated layer agrees with the straightforward expectations of the supernova hypothesis (Ellis, Fields, & Schramm 1996; Fields & Ellis 1999). These data greatly strengthen the case that supernova ejecta were deposited on the Earth about 3 Myr ago. This is a major new result, and one that finally gives a strong direct empirical basis for longstanding speculation and arguments based on indirect evidence regarding the possibility of near-Earth supernova explosions in the geological past.

In light of the Knie et al (2004) result, the time is ripe to take the supernova hypothesis seriously and investigate its consequences quantitatively. Specifically, in this paper we show how ferromanganese crusts can be used as telescopes to probe directly the r -process in supernovae.

2. Summary of the Data

Knie et al (2004) measured ^{60}Fe and ^{53}Mn (mean life $\tau_{53} = 5.3$ Myr) throughout the ferromanganese crust 237KD, using accelerator mass spectrometry techniques. Over the 28 layers studied, just 69 live ^{60}Fe atoms were detected. This signal was found uniquely in layers corresponding to a

time interval ¹

$$t = 2.4 - 3.2 \text{ Myr} \quad (1)$$

before the present. The resulting isotopic fraction is

$$\frac{{}^{60}\text{Fe}}{\text{Fe}} = (1.9 \pm 0.24) \times 10^{-15}, \quad (2)$$

which is a highly significant result, despite its small magnitude, testifying to the sensitivity and precision of the experiment. The principal uncertainty in this result is due to instrumental background, essentially the contamination due to stable isobars.

Given the crust size and Fe composition, the isotopic fraction can be used to infer a surface density of ${}^{60}\text{Fe}$. To do this, one must also correct for radioactive decay and geometric effects, and the result is

$$N_{60,\text{obs}} = 2.9 \times 10^6 \text{ atoms/cm}^2. \quad (3)$$

In order to infer the incident fluence of ${}^{60}\text{Fe}$, this result must be corrected for geometry and an uptake factor. Assuming the terrestrial fallout is isotropic, a geometric factor $1/4$ arises from the ratio of the Earth’s full cross section ($\pi R_{\oplus}^2$) to its total surface area ($4\pi R_{\oplus}^2$). Also, the crust incorporates only a certain percentage of all deposited material incident upon it, and a so-called ‘uptake factor’ $U < 1$ accounts for this effect (for further discussion, see Knie et al 2004; Knie et al. 1999), leading to

$$N_i(t) = \frac{U}{4} F_i \exp(-t/\tau). \quad (4)$$

The uptake factors are different for each sample and every element, and thus cannot be evaluated independently of the details of the sample. Consequently, in the following discussion we present results for $U = 1$, with the understanding that all the results presented must be reduced by the appropriate uptake factor, which may be inferred from studies of stable isotopes.

Knie et al (2004) use Mn and Fe data to infer an iron uptake factor $U_{\text{Fe}} \simeq 0.006$, from which they infer an incident fluence of

$$F_{60,\oplus} = 2.0 \times 10^9 \text{ atoms/cm}^2. \quad (5)$$

This signal encodes key information about the supernova explosion, as we now see.

3. Predictions: Supernova Distance and Other Radioisotope Signals

The deposited amount of supernova-produced material of an isotope i is dependent on the distance D of the event and the total ejected mass M_i . The fluence, i.e., the number of atoms per

¹The crust was analyzed with two different time resolutions but even the smaller timespan of 0.3 Myr in the finer resolution analysis is expected to be an artifact representing an upper limit on the physical deposition timescale, which is likely to be of order ~ 10 kyr.

unit area that arrive at the Earth, is (Fields & Ellis 1999; Ellis, Fields, & Schramm 1996):

$$F_i = \frac{M_{\text{ej},i}}{4\pi A_i m_p D^2}, \quad (6)$$

where A_i is the mass number of isotope i and m_p is the proton mass. Present-day measurements of the fluence must also be corrected for decay.

The isotopic fraction of i in the crust is determined by the relative amounts of supernova ejecta and substrate deposited within the observed time interval. If the substrate material has density ρ and deposition rate $\dot{h}$, then the total mass flux onto the crust is $j = \rho \dot{h}$, and the iron flux onto the crust is $X_{\text{Fe}} j$, where $X_{\text{Fe}} \sim 0.2$ (Hein et al. 2000) is the iron mass fraction in the crust, and other elements i can be treated in a corresponding way. Finally, if the supernova signal is measured with a time resolution Δt , the surface density of crust atoms deposited is $N = X_{\text{Fe}} j \Delta t / A m_p$, where $A \approx A_i$ is the mean mass number of the element, and so the isotopic fraction of i in the crust is

$$\frac{N_i}{N} = \frac{U_i M_{\text{ej},i}}{16\pi X_{\text{Fe}} \rho \dot{h} \Delta t D^2} e^{-t/\tau_i} \quad (7)$$

$$= 1.9 \times 10^{-15} \left(\frac{M_{\text{ej},i}}{2 \times 10^{-5} M_{\odot}} \right) \left(\frac{0.3 \text{ Myr}}{\Delta t} \right) \left(\frac{30 \text{ pc}}{D} \right)^2, \quad (8)$$

where the fiducial numbers are for the ^{60}Fe signal at 3 Myr, with $\rho = 5 \text{ g cm}^{-3}$ and $\dot{h} = 2.5 \text{ mm Myr}^{-1}$.

The confirmed observation of a live ^{60}Fe signal spurs us to predict the inventory of all long-lived radioisotopes. For each, given the yield $M_{\text{ej},j}$, one can readily compute the fluence and/or surface density, based on the ^{60}Fe signal:

$$N_j = \frac{U_j}{U_{\text{Fe}}} \frac{60}{A_j} \frac{M_{\text{ej},j}}{M_{\text{ej},60}} N_{60} e^{(1/\tau_{60} - 1/\tau_i)t}, \quad (9)$$

a result that is independent of supernova distance D . Consequently, detection of any radioisotopic species j would immediately measure the ratio of its yield relative to ^{60}Fe , and thus probe supernova nucleosynthesis - assuming that the uptake factors U are known, and that the Earth receives an isotropic ‘fair sample’ of all supernova ejecta ². Indeed, even a significant upper limit could provide important information.

The explosion signals divide into radioisotopes from three sources: supernovae, the r -process and cosmic rays, and we consider each of these in turn.

²The uptake factors U can be obtained from studies of stable isotopes or, in the case of ^{244}Pu , from studies of the recent deposition of anthropogenic Pu on the surface of the crust (Wallner et al. 2004).

3.1. Supernova Isotopes

To get an impression of the distance of the supernova, one needs to know the ^{60}Fe yield $M_{60\text{Fe}}$ of a supernova. This and the yields of other isotopes have been calculated in Woosley & Weaver (1995), and more recently again in Rauscher, Heger, Hoffman, & Woosley (2002). The 12 supernova explosion models of Woosley & Weaver (1995) span a mass range from $11 - 40M_{\odot}$, and the 5 newer calculations in Rauscher, Heger, Hoffman, & Woosley (2002) cover the range $15 - 25M_{\odot}$. The latter give a higher ^{60}Fe yield than those calculated in Woosley & Weaver (1995). A common feature of the ^{60}Fe yields in the two papers is strong variation over the observed mass range, which attests to the difficulty of calculating the production of this element, reflected also in differences between the results of Rauscher, Heger, Hoffman, & Woosley (2002) and Woosley & Weaver (1995) for supernovae of similar masses. Nevertheless, we use these mass yields together with eq. (6) to calculate the possible supernova distance D , albeit with considerable uncertainties induced by the different mass yields.

The results of this calculation can be seen in Fig. 1, where we see that the distances cover a range between ~ 10 pc and ~ 100 pc. This range is similar to that found in the similar analysis of Knie et al (2004). More generally, this distance range falls within limits set by Ellis & Schramm (1995) and Ellis, Fields, & Schramm (1996), who note that a supernova closer than 10 pc could be devastating to life on earth (e.g., Ruderman 1975), whereas a supernova that was too far away would not have the power to overcome the solar wind and deposit a noticeable amount of isotopes.

Having estimated the distance, it is now easy to obtain the surface densities (4) of other sufficiently long-lived supernova-produced elements like ^{36}Cl ($\tau = 0.435$ Myr), ^{53}Mn ($\tau = 4.5$ Myr) and ^{26}Al ($\tau = 1.03$ Myr), using again the values of Rauscher, Heger, Hoffman, & Woosley (2002) and Woosley & Weaver (1995) for the ejected masses of ^{36}Cl , ^{53}Mn and ^{26}Al . The results for ^{53}Mn and ^{26}Al are shown in Figure 2, and vary wildly, due to the variations in the ^{60}Fe mass yields. A better prediction of the ^{60}Fe yields would therefore be a valuable input into future calculations. On the other hand, the detection in the sample of Knie et al (2004) (or in comparable samples) of other isotopes, besides ^{60}Fe , whose yields are not as wildly varying (e.g. ^{26}Al), would allow for a more accurate calculation of the distance and hence stronger predictions for the surface densities of other supernova isotopes, and possibly the supernova mass as well. At the present stage, the results shown in Figure 2 can be used only to get a feel for the expected order of magnitude (without uptake factors) of the surface densities of supernova isotopes. As we see in the next Section, the corresponding results for ^{36}Cl are not worth showing because of the high expected background.

3.2. r -Process Elements

The r -process elements present a special case. These are nuclides which are created when ‘seed’ nuclei – typically iron-peak elements – are exposed to a rapid burst of neutrons. Whilst the microphysics and thermodynamical requirements for the r -process are fairly well understood,




Fig. 1.— The supernova distance implied by the Knie et al (2004) result, shown as a function of progenitor mass. Results are presented for the theoretical yields of Rauscher, Heger, Hoffman, & Woosley (2002) and Woosley & Weaver (1995). The curves scale with the ^{60}Fe yield as $M_{\text{ej},60}^{-1/2}$ (γ), and the large variations in distance reflect the large variations in the yields with progenitor mass, and the differences between the curves give a sense of the range of theoretical uncertainty in the yields.




Fig. 2.— Predicted terrestrial surface densities N_i as functions of supernova progenitor mass (eq. 6). For ^{53}Mn and ^{26}Al , the background levels are indicated. Results are corrected for geometry and decay, but are not corrected for site- and element-dependent uptake factors U which appear in eq. (4). As in Figure 1, variations in the predicted ^{60}Fe yields are the dominant cause of the large excursions with varying mass. (a) Results using Woosley & Weaver (1995) yields. (b) Results using the Rauscher, Heger, Hoffman, & Woosley (2002) yields.

its astrophysical site is not yet known; both core-collapse supernovae (e.g., Woosley et al. 1994) and neutron-star mergers (Eichler, Livio, Piran, & Schramm 1989) have been studied in some detail. Consequently, a firm correlated detection of a known supernova-produced isotope (e.g., ^{36}Cl , ^{53}Mn , ^{26}Al , or ^{60}Fe) with an r -process isotope (^{244}Pu , ^{146}Sm , or ^{182}Hf) would be of the utmost importance: it would provide direct proof that at least some supernovae do generate r -process elements. Intriguingly, Wallner et al. (2004) report the detection of a single live ^{244}Pu atom in the same crust that contained the original live ^{60}Fe signal (Knie et al. 1999). One obviously should not overinterpret such a marginal result, but we are encouraged that ^{244}Pu searches are becoming feasible, and we advocate strongly further searches for all r -process elements.

In order to get a feel for the expected order of magnitude of the surface density of ^{244}Pu , ^{146}Sm and ^{182}Hf , we must determine the supernova yields of these elements. We do this empirically, by noting that, in metal-poor Galactic halo stars, the heaviest elements are always distributed in the same ratios as the solar r -process component. This result implies that, at least for the heaviest elements (including the species of interest here), the r -process yield distribution is ‘universal’, i.e., always produced in the solar ratios: (Fields, Truran, & Cowan 2002) and refs. therein. Thus, all r -process isotopes are fixed once one is known. We make this normalization using the halo star observations, and choose Eu. Fields, Truran, & Cowan (2002) use observations of Eu and Fe in halo stars to deduce that one supernova produces $m_{\text{Eu}} \approx 5 \times 10^{-7} M_{\odot}$. It then follows that any other species r has $m_r = (A_r/152)(r/\text{Eu})_{\odot,r} m_{\text{Eu}}$. For solar abundances of the r -process elements, we use a recent theoretical model (Pfeiffer, Kratz, & Thielemann 1997; Cowan et al. 1999), from which we extract the following ratios: $(\text{Sm}/\text{Eu})_{\odot,r} = 1.1$, $(\text{Hf}/\text{Eu})_{\odot,r} = 1.1$, and $(\text{Pu}/\text{Eu})_{\odot,r} = 0.12$; this gives supernova yields

$$m_{\text{Sm}} = 5.4 \times 10^{-7} M_{\odot}, \quad m_{\text{Hf}} = 6.8 \times 10^{-7} M_{\odot}, \quad m_{\text{Pu}} = 0.88 \times 10^{-7} M_{\odot}. \quad (10)$$

Using eq. (4), one can now calculate the surface densities of the r -process elements in question, and our results are shown in Figure 2. Notice again the wild variations at the present stage, which allow the shown results only to give an impression of the expected orders of magnitude of the surface densities.

Estimates of the expected backgrounds are given in the next Section.

3.3. Cosmic-Ray Production: ^{10}Be

Finally, we consider the possibility of a supernova signal due to ^{10}Be by cosmic rays. The explosion of a nearby supernova enhances the cosmic ray flux at Earth, and the element that is dominantly produced is ^{10}Be ($\tau = 2.18$ Myr). Ellis, Fields, & Schramm (1996) estimate that an average supernova will put a fraction $\xi_{\text{SN}} \gtrsim 1/100$ of its mechanical energy E_{SN} into the production

of cosmic rays with an average energy of $\langle E_{CR} \rangle = 1$ GeV. Thus we get a cosmic ray fluence of

$$\Phi \Delta t = \xi_{CR} f_{CR} \frac{\langle E_{SN} \rangle}{4\pi D^2 \langle E_{CR} \rangle}, \quad (11)$$

The deflecting effects of the solar wind mean that only some fraction f_{CR} of cosmic rays penetrate to the Earth. Following Fields & Ellis (1999), we take $f_{CR} = 1/10$.

The enhanced cosmic-ray fluence at Earth increases radioisotope production in the atmosphere, according to

$$F_i = Y_i \Phi_p \Delta t, \quad (12)$$

where Y_i is the branching ratio for cosmic-ray production of isotope i ; a useful tabulation of these appears in O’Brien, de La Zerda Lerner, Shea, & Smart (1991). Using eq. (12) with $Y(^{10}\text{Be}) = 1.36 \times 10^{-2}$, Φ from eq. (11), and the distances from Figure 1, one finds a ^{10}Be fluence of at most $2.8 \times 10^8 \text{at/cm}^2$, and a surface density of at most $2.0 \times 10^7 \text{atoms cm}^{-2}$.

4. Backgrounds

We now consider the dominant sources of backgrounds to these various signals, which originate from different sources.

4.1. Background for ^{53}Mn and ^{26}Al : Meteoritic Infall

The backgrounds for ^{26}Al and ^{53}Mn are dominated by the infall of meteorites. These have been exposed to cosmic rays, causing an enhancement of certain elements due to spallation. An introductory discussion of meteoritic backgrounds can be found in Fields & Ellis (1999). The flux Φ_i of infalling isotopes can be calculated as

$$\Phi_i = \frac{J X_i}{4\pi R_\oplus^2 A_i m_p}, \quad (13)$$

where $J \approx 4 \times 10^{10} \text{g/yr}$ is the infalling meteoritic mass rate (Peucker-Ehrenbrink 1996; Esser & Turekian 1988; Love & Brownlee 1995) and X_i is the mass fraction of isotope i found in Solar-System meteorites. Michel et al. (1991) examine the production of isotopes in meteorites due to cosmic rays. They give their main tabulated results in terms of the specific activity Γ_i , which is the number of decays per minute and kg of iron or rock, respectively, for ^{53}Mn or ^{26}Al . We take the fiducial values of 400 dpm/kg Fe and 40 dpm/kg Si. In fact, ^{26}Al has more than one production branch, but as Si is the most dominant element we will use it for an estimate of the background. We assume the infalling meteorites to have an iron mass fraction $X_{Fe} = 0.19$ and a silicon mass fraction $X_{Si} = 0.25$ (Anders & Grevesse 1989). For a meteorite with such mass fractions, we obtain

$X_i = m_i \tau_i \Gamma_i X_{Fe/Si}$ and therefore $X(^{53}\text{Mn}) = 1.9 \times 10^{-11}$ and $X(^{26}\text{Al}) = 2.4 \times 10^{-13}$. With these, we find fluxes of

$$\Phi(^{53}\text{Mn}) = 1.7 \times 10^9 \text{ Myr}^{-1} \text{ cm}^{-2}, \quad (14)$$

$$\Phi(^{26}\text{Al}) = 4.3 \times 10^7 \text{ Myr}^{-1} \text{ cm}^{-2}. \quad (15)$$

In order to calculate the background fluence and surface density we have to multiply the flux by the time resolution. Recalling that the ^{60}Fe signal in Knie et al (2004) spreads over a time of approximately $\Delta t \simeq 1/3$ Myr, we obtain surface densities of

$$N(^{53}\text{Mn}) = 8.4 \times 10^7 \text{ atoms cm}^{-2}, \quad (16)$$

$$N(^{26}\text{Al}) = 2.4 \times 10^5 \text{ atoms cm}^{-2}. \quad (17)$$

In comparison with the signal strengths of order $10^8 - 10^9 \text{ atoms cm}^{-2}$ for ^{53}Mn and $10^7 - 10^8 \text{ atoms cm}^{-2}$ for ^{26}Al , these backgrounds are negligible.

4.2. Background for ^{10}Be and ^{36}Cl : Cosmic-Ray Interactions

A different approach has to be taken for calculating the backgrounds to ^{36}Cl and ^{10}Be , as these nuclides are primarily produced by the direct interaction of cosmic rays with the atmosphere. Using eq. (12), with the ambient cosmic-ray proton flux $\Phi_p \simeq 10 \text{ cm}^{-2} \text{ s}^{-1}$ and a time resolution $\Delta t = 0.3$ Myr, for ^{36}Cl one finds $Y(^{36}\text{Cl}) = 4.74 \times 10^{-4}$. Consequently, one obtains a present-day surface density of

$$N(^{36}\text{Cl}) = 2.0 \times 10^7 \text{ atoms cm}^{-2}. \quad (18)$$

This is to be compared with a signal strength $\sim 10^4 - 10^6 \text{ atoms cm}^{-2}$ and we therefore do not expect the detection of ^{36}Cl in the ferromanganese crust sample of Knie et al (2004) or in similar samples.

4.3. Background for ^{244}Pu , ^{146}Sm , and ^{182}Hf : Fission

As ^{146}Sm , ^{182}Hf and ^{244}Pu are extinct radioactive isotopes, they do not have a significant abundance in Solar-System meteorites. Moreover, they are not produced in sufficient amounts by cosmic ray spallation. The only background source is spontaneous fission of actinides. Such decays lead to a distribution of daughter products, but these only contribute significantly to ^{146}Sm , not to ^{182}Hf .

We estimate the spontaneous fission background of ^{146}Sm as follows. The dominant parent nucleus is ^{238}U , with a fission lifetime $\tau_{238,\text{sf}} = 1.2 \times 10^{16}$ yr. If the fission branching ratio to ^{146}Sm is $f \sim 0.1$, then the equilibrium ^{146}Sm fission background thus has $(^{146}\text{Sm}/^{238}\text{U})_{\text{eq}} = f \tau_{146}/\tau_{238,\text{sf}}$. The ^{146}Sm background flux onto the crust is then $\Phi_{146} = (^{146}\text{Sm}/^{238}\text{U})_{\text{eq}} X_{238j}/238m_p$ (where j is

the total mass flux onto the sample, and $X_{238} \sim 3 \times 10^{-6}$ is the mass fraction of ^{238}U in the Earth’s crust). Thus the background accumulated in a resolution time $\Delta t \sim 0.3$ Myr is

$$N_{146,\text{bgd}} = U f \frac{\tau_{146}}{\tau_{238,\text{sf}}} X_{238} \frac{j \Delta t}{238 m_p} \simeq 3.6 \times 10^7 U \quad (19)$$

which is larger than our prediction by about an order of magnitude. Thus we conclude that the ^{146}Sm is probably too small to see.

Since ^{244}Pu is *heavier* than ambient U and Th, it is not produced by fission. Apart from recent human activity, the only known possible source of ^{244}Pu is the r -process. Moreover, anthropogenic ^{244}Pu is expected to be negligible in the ferromanganese crusts of interest.

5. Discussion

Following the prediction of Ellis, Fields, & Schramm (1996), Knie et al. (1999) found initial evidence for supernova ^{60}Fe in a ferromanganese crust, prompting Fields & Ellis (1999) to discuss other isotopic signatures of a nearby supernova. The exciting new result of Knie et al (2004) confirms the earlier observation with higher significance and more precise dating, motivating this paper. Using the data of Knie et al (2004) to estimate the local interstellar fluence of ^{60}Fe and combining this with the total mass of ^{60}Fe ejected by diverse supernovae (Rauscher, Heger, Hoffman, & Woosley 2002; Woosley & Weaver 1995), we have inferred a supernova distance range $D \sim 10\text{--}100$ pc, consistent with Knie et al (2004).

The ^{60}Fe detection and nucleosynthesis models together allow us to make distance-independent predictions for other radioisotopes. For supernova-produced isotopes such as ^{26}Al , ^{53}Mn and ^{36}Cl , we find that yields of the first two of these isotopes are likely to be detectable above background. We also consider the r -process elements ^{244}Pu , ^{182}Hf , and ^{146}Sm , finding again that the first two are likely to exceed background. If the supernova responsible for the ^{60}Fe observations of Knie et al (2004) was also a site for the r -process, clear signals should be visible. This would finally provide direct proof that supernovae are sites for the r -process.

The firm detection of ^{60}Fe also adds support to recent ideas regarding the origin of the Local Bubble. Smith & Cox (2001) concluded that several events are needed to create the Local Bubble, the last occurring within the past 5 Myr in order to maintain the high interstellar temperature. Maíz-Apellániz (2001) argued that the need for multiple events points to correlated explosions within an OB association, and specifically suggested the Lower Centarus Crux (LCC) of the Sco-Cen association as the likely site. Benítez, Maíz-Apellániz, & Canelles (2002) have examined the possibility that this OB association could have been the site of the ^{60}Fe progenitor. From kinematic considerations they find that the closest approach of the LCC was about 100 pc, with considerable uncertainties. Taken at face value, this result is in the upper part of our range of distance estimates. This suggests that either the event was a distant one (and thus the ^{60}Fe yield was high), or that the LCC approach was closer than estimated, or that the LCC was not the event site.

The accuracies of our predictions for the fluences of these isotopes are limited by the wide variations in the ^{60}Fe yields found in calculations for different progenitor masses, and between different model calculations (Rauscher, Heger, Hoffman, & Woosley 2002; Woosley & Weaver 1995). Therefore, at the present stage our results give only approximate estimates of the expected orders of magnitude of the fluences of supernova isotopes. This underlines the great need for more theoretical studies of ^{60}Fe production in supernovae, able to reduce the calculational uncertainties. Such a reduction would be necessary before one could predict any biological effects that might be associated with the supernova responsible for the ^{60}Fe observations of Knie et al. (1999) and Knie et al (2004).

Measurements using other ferromanganese crusts would be most welcome, and observations of other supernova radioisotopes have high priority. These would enable one to evade the uncertainties associated with the ^{60}Fe yield. Another helpful calibration point comes from the recent evidence (Smith 2004) for the γ -ray line signal from the decay of interstellar ^{60}Fe . This reflects the current Galactic ^{60}Fe inventory, and constrains ^{60}Fe nucleosynthesis averaged over all sources (Prantzos 2004).

The important results of Knie et al. (1999) and Knie et al (2004) open up a new era of ‘deep-ocean crust astronomy’. After providing already the first observations of the terrestrial impact of a nearby supernova, their measurements now offer the prospect of verifying directly that the r -process takes place, at least partly, in ejecta from supernovae.

We thank Günther Korschinek and Klaus Knie for invaluable discussions regarding their ^{60}Fe data. This material is based upon work supported by the National Science Foundation under Grant No. AST-0092939

REFERENCES

- Anders, E. & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
- Benítez, N., Maíz-Apellániz, J., & Canelles, M. 2002, *Physical Review Letters*, 88, 081101
- Cowan, J. J., et al 1999, *ApJ*, 521, 194
- Eichler, D., Livio, M., Piran, T., & Schramm, D. N. 1989, *Nature*, 340, 126
- Ellis, J., Fields, B. D., & Schramm, D. N. 1996, *ApJ*, 470, 1227
- Ellis, J. & Schramm, D. N. 1995, *Proc. Nat. Acad. Sci.*, 92, 235
- Esser, B. K. & Turekian, K. K. 1988, *Geochim. Cosmochim. Acta*, 52, 1383
- Fields, B. D. & Ellis, J. 1999, *New Astronomy*, 4, 419
- Fields, B. D., Truran, J. W., & Cowan, J. J. 2002, *ApJ*, 575, 845

- Frisch, P. C. 1995, *Space Science Reviews*, 72, 499
- Hein, J.R., et al. 2000, in *Handbook of Marine Mineral Deposits*, ed. D.S. Cronan (New York: CRC Press), 239
- Knie, K., et al. 1999, *Phys. Rev. Lett.*, 83, 18
- Knie, K. et al. 2004, *Phys. Rev. Lett.*, 93, 171103
- Love, S. G. & Brownlee, D. E. 1995, *Science*, 262, 550
- Maíz-Apellániz, J. 2001, *ApJ*, 560, L83
- McCullough, P. R., Fields, B. D., & Pavlidou, V. 2002, *ApJ*, 576, L41
- Michel, R., Dragovitsch, P., Cloth, P., Dagge, G., & Filges, D. 1991, *Meteoritics*, 26, 221
- O’Brien, K., de La Zerda Lerner, A., Shea, M. A., & Smart, D. F. 1991, *The Sun in Time*, 317
- Peucker-Ehrenbrink, B. 1996, *Geochim. Cosmochim. Acta*, 60, 3187
- Pfeiffer, B., Kratz, K.-L., & Thielemann, F.-K. 1997, *Zeitschrift fur Physik A Hadrons and Nuclei*, 357, 235
- Prantzos, N. 2004, *A&A*, 420, 1033
- Rauscher, T., Heger, A., Hoffman, R. D., & Woosley, S. E. 2002, *ApJ*, 576, 323
- Ruderman, M. 1975, *Science*, 184, 1079
- Smith, D. M. 2004, *New Astronomy Review*, 48, 87
- Smith, R. K. & Cox, D. P. 2001, *ApJS*, 134, 283
- Wallner, C., Faestermann, T., Gerstmann, U., Knie, K., Korschinek, G., Lierse, C., & Rugel, G. 2004, *New Astronomy Review*, 48, 145
- Woosley, S. E. & Weaver, T. A. 1995, *ApJS*, 101, 181
- Woosley, S. E., Wilson, J. R., Mathews, G. J., Hoffman, R. D., & Meyer, B. S. 1994, *ApJ*, 433, 229